ORTHOINTSPINESMUSCLE

Troy Roberson, MD 8450 Northwest Blvd. Indianapolis, IN 46278 317.802.2477 • Ortholndy.com

Open or Arthroscopic HAGL and/or Subscapularis Repair

Name: ____

Date: _____ /____ /____

Phase 0: Quiet (Weeks 1 to 3)

- Regular sling for six weeks
- No shoulder rehab
- No pendulums
- Elbow and hand only

Phase I: Passive (Weeks 4 to 5)

- Pendulums to warm-up
- Passive range of motion
- Start supine external rotation 0 degrees, then gradually progress, but limit ER to 30 degrees
- Supine forward elevation 90 degrees
- No internal rotation

Phase II: Active (Weeks 6 to 10)

- Internal rotation to belt line gradually increase to full
- Full elevation and gradually increase ER limit ER progression to 30 degrees
- Active range of motion with passive stretch to prescribed limits
- Supine to start external rotation gradually increase to full
- Supine to start forward elevation full progress to seated
- Internal rotation gradually increase to full

Phase III: Resisted (Week 10)

- Pendulums to warm-up and continue with Phase II
- Scapular mobilization
- External and internal rotation
- Standing forward punch
- Scapular mobilization seated rows and shoulder shrugs

Weight Training (Week 12) *select patients

- Keep hands within eyesight and elbows bent
- Minimize overhead activities
- No military press, pull-down behind head or wide grip bench

Return to Activities

- Computer: 4 weeks
- Golf: 4 months
- Tennis: 5 months