

Troy Roberson, MD 8450 Northwest Blvd. Indianapolis, IN 46278 317.802.2477 • Ortholndy.com

Rotator Cuff	Repair	(Less	than	3cm)
--------------	--------	-------	------	------

Name:	D	/	/
Name:	l)ate:	, ,	/
Name	_ Date/	/	

Phase 0: Quiet (Weeks 0 to 1)

- Quiet in sling with elbow/wrist/hand
- UltraSling for weeks 0 to 3
- Begin active scapular retraction/protraction exercises with therapist cueing

Phase I: Passive (Weeks 2 to 6)

- Pendulums to warm-up beginning week one
- Supine external rotation 0 to 30 degrees beginning at two weeks with progression to full PROM by six weeks
- Supine forward elevation 0 to 90 degrees beginning at two weeks with progression to full PROM by six weeks
- Progress to upright as tolerated with ER and FE

Phase II: Active (Weeks 7 to 9)

- Pendulums to warm-up
- Active range of motion with **terminal stretch**
- Supine → external rotation after six weeks; progress gradually to full
- Supine → forward elevation after six weeks; progress **gradually** to full
- Begin active biceps
- Internal rotation full (begin behind the back)
- Begin AROM in supine and progress to upright

Phase III: Resisted (Week 10)

- Pendulums to warm up and continue with Phase II
- External and internal rotation
- Standing forward punch
- Seated rows
- Shoulder shrugs and biceps curls

Weight Training (Week 12)

- Keep hands within eyesight, keep elbows bent, no long lever arms
- Minimize overhead activities (below shoulder)
- No Military press, pull-down behind head or wide grip bench

Initiation of Interval Sport Programs

Golf: 3 monthsTennis: 4 monthsSki: 3 to 4 months