

Scott Gudeman, MD 1260 Innovation Pkwy., Suite 100 Greenwood, IN 46143 317.884.5161 • Ortholndy.com ScottGudemanMD.com

Subacromial Decompression, Arthroscopic Debridement, Acromioplasty, Distal Clavical Excision, SLAP I Debridment Rehabilitation Protocol

Days 1 to 14

- Office visit to have a dressing changed, one to two days post-op
- Cryotherapy as needed to control pain and swelling
- Patient should be wearing a sling at all times except when performing exercises
- Suture removal office visit, seven days post-op

Goals

- Restore non-painful range of motion
- Retard muscular atrophy
- Decrease pain and inflammation
- Improve postural awareness

Exercises

- Begin physical therapy soon after surgery
- Passive range of motion
- Active assertive range of motion
- Grip strength
- Elbow/wrist/hand active range of motion
- Pendulum exercises/Codman's
- Pulley/canes
- Scapular stabilization to begin at day seven
- Electronic muscle stimulation as needed
- Pain level should guide exercises and activity
- Upper body exercises

Weeks 2 to 4

- Continue with cryotherapy as needed
- Patient should be out of sling at week four

Exercises

- Focus on active range of motion
- Begin rotator cuff strengthening as tolerated
- Isometrics
- Light theraband exercises
- Stretching
- Continue with scapular stabilization exercises

1 ______ Rev. 10/15

Week Four to Eight

Goals

- Return to office visit for third post-op visit
- Continue physical therapy as needed to progress strengthening: active range of motion
- Full range of motion at four weeks
- No pain or swelling
- Regain and improve muscular strength

Exercises

- Continue all strengthening, progress as tolerable
- Isotonic strengthening (theraband, dumbbells)
- Proprioceptive neuromuscular facilitation
- Strengthen scapulothoracic musculature
- Electronic muscle stimulation
- Passive stretching at end ranges to maintain flexibility
- UT mobs and scapular mobs

Weeks 8 to 12

Exercises

- Advance strengthening as tolerated
- Eccentrically resisted motions
- Closed kinetic chain

Months Two to Three

• Functional progression back to sport/work

Helping you achieve the optimal activity level for your lifestyle is my first priority.

- Scott Gudeman, MD



Scan here to visit ScottGudemanMD.com.